

UC Irvine

UC Irvine Previously Published Works

Title

Inhibition of Intestinal Thiamin Transport in Rat Model of Sepsis.

Permalink

<https://escholarship.org/uc/item/6284j6v8>

Journal

Critical care medicine, 44(9)

ISSN

0090-3493

Authors

Sassoon, Catherine S
Zhu, Ercheng
Fang, Liwei
et al.

Publication Date

2016-09-01

DOI

10.1097/ccm.0000000000001745

Peer reviewed

Inhibition of Intestinal Thiamin Transport in Rat Model of Sepsis

Catherine S. Sassoon, MD^{1,2}; Ercheng Zhu, PhD^{1,2}; Liwei Fang, BSc¹; Veedamali S. Subramanian, PhD^{2,3}; Hamid M. Said, PhD^{2,3}

Objectives: Thiamin deficiency is highly prevalent in patients with sepsis, but the mechanism by which sepsis induces thiamin deficiency is unknown. This study aimed to determine the influence of various severity of sepsis on carrier-mediated intestinal thiamin uptake, level of expressions of thiamin transporters (thiamin transporter-1 and thiamin transporter-2), and mitochondrial thiamin pyrophosphate transporter.

Design: Randomized controlled study.

Setting: Research laboratory at a Veterans Affairs Medical Center.

Subjects: Twenty-four Sprague-Dawley rats were randomized into controls, mild, moderate, and severe sepsis with equal number of animals in each group.

Interventions: Sepsis was induced by cecal ligation and puncture with the cecum ligated below the cecal valve at 25%, 50%, and 75% of cecal length, defined as severe, moderate, and mild sepsis, respectively. Control animals underwent laparotomy only.

Measurements and Main Results: After 2 days of induced sepsis, carrier-mediated intestinal thiamin uptake was measured using [³H]thiamin. Expressions of thiamin transporter-1, thiamin transporter-2, and mitochondrial thiamin pyrophosphate transporter proteins and messenger RNA were measured. Proinflammatory

cytokines (interleukin-1 β and interleukin-6) and adenosine triphosphate were also measured. Sepsis inhibited [³H]thiamin uptake, and the inhibition was a function of sepsis severity. Both cell membrane thiamin transporters and mitochondrial thiamin pyrophosphate transporter expression levels were suppressed; also levels of adenosine triphosphate in the intestine of animals with moderate and severe sepsis were significantly lower than that of sham-operated controls.

Conclusions: For the first time, we demonstrated that sepsis inhibited carrier-mediated intestinal thiamin uptake as a function of sepsis severity, suppressed thiamin transporters and mitochondrial thiamin pyrophosphate transporter, leading to adenosine triphosphate depletion. (*Crit Care Med* 2016; XX:00–00)

Key Words: intestinal thiamin uptake; mitochondria; sepsis; thiamin deficiency; thiamin pyrophosphate; thiamin transporter

The active form of thiamin, thiamin pyrophosphate, represents ~85–90% of total cellular thiamin, and is generated exclusively in the cytoplasm and transported into the mitochondria for utilization of cellular metabolic process. Thiamin pyrophosphate is indispensable for normal mitochondrial function and structural integrity due to its involvement as a cofactor in oxidative energy metabolism, adenosine triphosphate (ATP) production, and maintenance of cellular redox state by reducing cellular oxidative stress (1). Thus, low intracellular levels of thiamin leads to impairment in oxidative energy metabolism (acute energy failure), a propensity for oxidative stress, and eventually apoptosis.

In critically ill adult and pediatric patients, thiamin deficiency is highly prevalent, ranged from 20% to 70% (2–4). In one retrospective study (5), the mortality rate of critically ill patients with thiamin deficiency was 72% compared with 44% without the deficiency. Thiamin deficiency appears to contribute to the high mortality, yet the risk factor(s) contributing to the deficiency were not investigated. In critically ill patients with sepsis, sepsis has been identified as an important risk factor for thiamin deficiency (2–4). However, the extent by which sepsis leads to thiamin deficiency is unknown. Thiamin is not synthesized endogenously, but has to be obtained exogenously

¹Department of Medicine, VA Long Beach Healthcare System, Long Beach, CA.

²Department of Medicine, University of California, Irvine, CA.

³Department of Physiology and Biophysics, University of California, Irvine, CA.

This work was conducted in the Department of Research and Development, VA Long Beach Healthcare System, Long Beach, CA.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website (<http://journals.lww.com/ccmjournal>).

Supported, in part, by the Department of Veterans Affairs Medical Research Service awarded to Drs. Sassoon and Said, and National Institutes of Health (DK-56061, DK-58057, AA-018071) awarded to Dr. Said.

Dr. Sassoon is a member of the Board of Directors of Barlow Respiratory Research in Los Angeles, CA (without compensation). The remaining authors have disclosed that they do not have any potential conflicts of interest.

For information regarding this article, E-mail: csassoon@uci.edu

Copyright © 2016 by the Society of Critical Care Medicine and Wolters Kluwer Health, Inc. All Rights Reserved.

DOI: 10.1097/CCM.0000000000001745

from nutrient via transport across the intestinal cell membrane, or from the circulation via transport to other organs. The cell membrane thiamin transporters (THTRs), THTR-1 (encoded in *SLC19A2* gene) and THTR-2 (encoded in *SLC19A3* gene), mediate intestinal thiamin transport process, whereas the uptake of thiamin pyrophosphate into the mitochondria occurs via a carrier-mediated mechanism that involves the mitochondrial thiamin pyrophosphate transporter (MTPPT, encoded in *SLC25A19* gene) (6). The effect of sepsis on intestinal thiamin uptake, and its influence on intestinal THTRs, THTR-1 and THTR-2, and MTPPT is unknown. We hypothesize that sepsis produces thiamin deficiency, at least in part, by inhibition of thiamin uptake via suppression of intestinal THTRs. Furthermore, sepsis suppresses MTPPT resulting in reduced ATP generation. The objective of this study is to investigate in experimental model of sepsis in rats, the influence of sepsis and degree of severity on intestinal thiamin uptake, and the state of THTRs in cytoplasm and mitochondria.

MATERIAL AND METHODS

Subjects, Surgical Procedures, Animal Monitoring and Care

This study received approval from the Research and Development Subcommittee on Animal Studies of the Veterans Affairs Long Beach Healthcare System. We studied a total of 24 male pathogen-free Sprague-Dawley rats randomized into groups of sham control, severe, moderate, and mild degree of sepsis with equal number of animals in each group. Sepsis was produced using the cecal ligation and puncture (CLP) technique (7). The severity of sepsis was based on the extent of cecal ligation (see below) (7). Surgeries were performed under general anesthesia with an intramuscular mixture of ketamine hydrochloride 100 mg/kg and xylazine 5 mg/kg. A 2-cm midline incision of the skin and peritoneum was made in the lower abdomen. The cecum was identified, withdrawn through the incision, and ligated below the cecal valve at 25%, 50%, and 75% of the cecal length with a 3-0 silk suture to define severe, moderate, and mild degree of sepsis, respectively. Using an 18-gauge needle, the cecum was perforated in two locations on the antimesenteric surface and was gently compressed until feces were extruded to ensure patency of the holes. The sham control animals underwent similar procedures without the CLP. During surgery, rectal temperature was monitored, and maintained at 37°C using homeothermic warming pad (Right-Temp, Kent Scientific, Torrington, CT) and a heating lamp. Blood pressure was monitored noninvasively using the tail cuff method (CODA Monitor, Kent Scientific, Tomington, CT). Respiration, heart rate, and arterial oxygen saturation (SpO_2) were monitored and displayed in real time using pulse oximeter (MouseSTAT, Kent Scientific). The animals were allowed to recover for 2 days. Postoperatively, rectal temperature, blood pressure, respiration, and heart rate were measured twice a day. Antibiotic (gentamicin 0.5 mg/kg IM bid), and analgesic (buprenorphine 0.05 mg/kg subcutaneously every 6 hr) were administered immediately following surgery and for 2 days. Prewarmed 0.9% NaCl at

50 mL/kg SC was administered bid and as needed to maintain mean blood pressure of 70 mm Hg. Liquid acetaminophen 0.1 mg/mL was mixed in the drinking water. Softened rat chow was provided liberally; however, when oral intake was poor, animals were hand-fed twice a day with Ensure Complete 10 mL/kg (Abbott Nutrition, Columbus, OH). Animals that died, mostly because of hypotension, were replaced to maintain equal number of animals in each group ($n = 6$).

Measurement of Thiamin Uptake

After 2 days of sepsis, under general anesthesia, animals underwent laparotomy, and thiamin uptake measurement in the proximal jejunum using [^3H]thiamin. In vivo, two intact jejunal loops (2 cm each in length) separated with another jejunal loop of the same length were prepared by suture ligation. In the first segment, 250 μL of Krebs-Ringer buffer containing [^3H]thiamin was injected intraluminally, and in the second segment, labeled and unlabeled thiamin mixed with the same incubation medium was injected.

Uptake of [^3H]thiamin (250 μM) by the carrier-mediated process was determined by subtracting uptake in the presence of a high pharmacologic concentration of unlabeled thiamin (1,000 μM) from uptake in its absence. Radioactivity uptake counting was measured after 5 minutes with Beckman Counter LS 6500 (Beckman Coulter, Fullerton, CA) (linear phase of uptake). Uptake data were expressed in fmol/mg tissue wet weight/5 min. All thiamin uptake experiments in septic rats were run simultaneously with weight-matched control rats. Following thiamin uptake measurements, blood sample was withdrawn from inferior vena cava for thiamin and cytokines determination. Rats were then euthanatized with pentobarbital 200 mg/kg, interperitoneally.

Determination of Thiamin Blood Concentration and Level of Serum Cytokines

Whole blood thiamin concentration was measured using a microbiological test kit with *Lactobacillus fermentum*-coated microtiter plate (ID-Vit Vitamin B1 KIF001; Institut für Produktqualität GmbH, Berlin, Germany). Mean reference value of whole blood thiamin was $48.1 \pm 8.9 \mu\text{g/L}$ ($\pm \text{SD}$). Serum cytokines (interleukin [IL]-1 β and IL-6) were measured using enzyme-linked immunosorbent assay kit (Thermo Fisher Scientific, Grand Island, NY).

Determination of ATP

ATP in the jejunal tissue was measured colorimetrically using ATP assay Kit (ab83355; Abcam, Cambridge, MA) (8). Following dissection, tissue was flash frozen by immediate immersion in 100% isopentane (Mallinckrodt Baker, Phillipsburg, NJ) prechilled on dry ice. After tissue homogenization, samples were then deproteinated and neutralized by adding ice-cold Neutralization Solution (ab93299; Abcam, Cambridge, MA) at 1:24 (v:v), incubating on ice for 5 minutes, centrifuging at 10,000g for 2 minutes, and mixing with ice-cold ATP Assay Buffer at 1:5 (v:v). ATP standard curve was created, and ATP concentration in the tissue lysed from either controls or septic

animals was calculated relative to the standard curve, and normalized for protein concentration.

Determination of THTR-1, THTR-2, and MTPPT Protein and Messenger RNA (mRNA)

Frozen intestine tissues were processed for determination of THTR-1, THTR-2, and MTPPT protein abundance using standard western blotting. The intestine (40 mg) from the septic rats and paired controls was homogenized in protein buffer (T-PER78510; Thermo Scientific, Rockford, IL) and denatured for THTR-1 and THTR-2 protein analysis. Mitochondrial protein was extracted for MTPPT protein measurement. We employed Ponceau S as reference protein (9). The immunoreactive bands were analyzed using ImageQuant software, version 5.0 (Molecular Dynamics, Sunnyvale, CA).

For mRNA extraction, approximately 30 mg frozen intestine tissue was homogenized in TRIzol Reagent (Invitrogen, Carlsbad, CA) and total RNA was isolated according to the manufacturer's protocol. Following reverse transcription, the mRNA expression level was quantified in a real-time PCR system (CFX384 Touch Bio-Rad Laboratories, Hercules, CA) using SYBR Green Super mix (Bio-Rad Laboratories), and was performed in triplicate. 18S was used as the housekeeping gene. Specific primers for rats are as follows: THTR-1: forward 5'-GATGCTCCTACGTACTGCCC-3' and reverse 5'-GCAGGTAGGGAGTGAGGAAC-3'; THTR-2: forward 5'-TGATACTCTGCTTGTCGG-3' and reverse 5'-GTAAGAGTACGTCCAAACAG-3'; MTPPT: forward 5'-GGCCATACGCACCATG-3' and reverse 5'-GGGTCTTGCTGATGACTC-3', and 18S: forward 5'-GCAGAAATCCCACTCCCGACCC-3' and reverse 5'-CCCAAGCTCCAACTACGAGC-3'. Data were normalized to 18S and calculated using a relative relationship method supplied by the manufacturer (Bio-Rad Laboratories).

Statistical Analysis

Group size was determined from thiamin uptake values from the initial three animals based on minimum differences in a

mean of 0.30 and expected SD of residuals of 0.15. We required six animals per group to achieve a power of 0.80 and a α equal to 0.05 (analysis of variance [ANOVA]; SigmaPlot, version 12.0; SPSS, Chicago, IL). Values were expressed as mean \pm SE unless specifically indicated. A two-way ANOVA was used to compare physiologic variables among groups using the appropriate grouping variables of time (baseline and day 2) and severity of sepsis (mild, moderate, and severe as defined above). For other variables, one-way ANOVA was used for comparison among groups. When the f value was significant, post hoc analysis was performed using the Tukey test for pairwise multiple comparisons. Group differences were considered significant at p value less than or equal to 0.05.

RESULTS

The sepsis model employed in the present study resulted in animal survival rate of 100%, 90%, 86%, and 75% in sham control, mild, moderate, and severe sepsis groups, respectively. At baseline, compared with controls, body weight (BW), mean blood pressure, heart rate, respiratory rate, and SpO_2 were similar among groups, except body temperature was higher in all of the sepsis groups (Table S1, Supplemental Digital Content 1, <http://links.lww.com/CCM/B829>). After 2 days, BW tended to decrease in moderate and severe sepsis. Body temperature returned to levels that of controls, likely related to antibiotic administration. Hemodynamic, respiratory, and oxygenation variables were stable. The degree of proinflammatory cytokines (IL-1 β and IL-6) production is a function of sepsis severity (Table 1). Thiamin blood concentration in the septic animals was not significantly different from controls (Table 1).

Carrier-mediated thiamin uptake was reduced significantly in proportion to the severity of sepsis (Fig. 1). This was further supported by the significant correlation coefficient between thiamin uptake and IL-1 β (r^2 , 0.58; $p < 0.001$) and IL-6 (r^2 , 0.67; $p < 0.001$) (Fig. S1, Supplemental Digital Content 1, <http://links.lww.com/CCM/B829>). As thiamin pyrophosphate

TABLE 1. Serum Cytokines, Whole Blood Thiamin Concentration, and Intestine Adenosine Triphosphate in Control and Septic Animals

Variables	Control	Sepsis		
		Mild	Moderate	Severe
IL-1 β (pg/mL)	5.5 \pm 1.4	38.4 \pm 2.9	47.2 \pm 6.6	148.9 \pm 24.7 ^a
IL-6 (pg/mL)	11.8 \pm 4.6	78.4 \pm 15.8	143.8 \pm 12.8 ^b	465.1 \pm 46.2 ^a
Thiamin (μ g/L) ^c	51.9 \pm 1.1	52.5 \pm 0.8	51.1 \pm 1.3	50.6 \pm 0.3
Adenosine triphosphate (nmol/mg protein)	12.9 \pm 1.4	10.9 \pm 1.1	6.2 \pm 0.4 ^{d,e}	5.4 \pm 0.4 ^{d,f}

IL = interleukin.

^a $p < 0.001$ severe sepsis compared with control, mild, and moderate sepsis.

^b $p < 0.008$ moderate sepsis compared with control.

^cThiamin blood concentration $n = 4$.

^d $p < 0.001$ severe and moderate sepsis compared with control.

^e $p = 0.01$ moderate compared with mild sepsis.

^f $p = 0.003$ severe compared with mild sepsis.

Values are mean \pm SE; $n = 6$.




Figure 1. Intestinal carrier-mediated thiamin uptake. Thiamin uptake expressed as percent of controls. Values are mean \pm SE; $n = 6$ for each group. * $p < 0.001$: severe compared with mild sepsis and control; moderate compared with mild sepsis and control. † $p = 0.003$: severe compared with moderate sepsis. # $p = 0.001$: mild sepsis compared with control.

is essential for the conversion of pyruvate to acetyl-CoA in the generation of ATP, the reduced intestinal thiamin uptake led to a significantly decreased ATP in the moderate and severe sepsis groups, by 52% and 58%, respectively (Table 1).

The uptake of thiamin in the intestine is an active process mediated via cell membrane THTR-1 and THTR-2. As demonstrated in **Figure 2**, sepsis decreased both THTR-1 and

THTR-2 proteins significantly. Similarly, sepsis suppressed both THTR-1 and THTR-2 mRNA expressions (**Fig. 3**). The active form of thiamin, thiamin pyrophosphate, also requires a carrier-mediated transport for entry into the mitochondria. As shown in **Figure 4**, sepsis significantly reduced MTPPT protein and suppressed its mRNA expressions.

DISCUSSION

Sepsis is a systemic inflammatory response to infection with a prevalence for severe sepsis of 25–44% and a mortality rate of approximately 30% or greater (10). Few observational studies in critically ill patients have identified sepsis (2–4) as an important risk factor contributing to thiamin deficiency, and that the combined sepsis and thiamin deficiency was associated with higher mortality than sepsis alone (3). The present study is the first to demonstrate that 1) sepsis inhibited intestinal carrier-mediated thiamin uptake, and the inhibition was in proportion to the severity of sepsis (**Fig. 1**); 2) both THTR proteins and mRNA expression levels were suppressed (**Figs. 2 and 3**, respectively); and most importantly, 3) MTPPT protein and mRNA expressions were markedly decreased and associated with proportionate reduction in ATP in animals with moderate and severe sepsis.

Our animal model of sepsis mimicked human sepsis. The mortality rate of animals with severe sepsis was 25%, close to that reported mortality in human with sepsis (10). In contrast to a similar rat model of severe sepsis of 2 days duration, the present study had a significantly lower mortality rate of 25% compared with 70% as previously reported after 2 days of induced sepsis, 25% versus 70%, respectively (7). The reduced




Figure 2. Thiamin transporter (THTR)-1 (**A**) and THTR-2 (**B**) proteins. Values are mean \pm SE; $n = 4$ for each group. Representative gel images of respective proteins of different groups were obtained from the same gel. THTR-1: * $p = 0.002$ severe compared with control. * $p = 0.004$ moderate sepsis compared with control. ‡ $p = 0.027$ mild sepsis compared with control. THTR-2: * $p = 0.002$ severe sepsis compared with control. * $p = 0.006$ moderate sepsis compared with control. ‡ $p = 0.047$ mild sepsis compared with control.




Figure 3. Thiamin transporter-1 (A) and thiamin transporter-2 (B) messenger RNA (mRNA). Values are mean \pm SE; $n = 4$ for each group. Thiamin transporter-1: * $p < 0.001$ severe and moderate sepsis compared with control. * $p = 0.001$ mild sepsis compared with control. ‡ $p = 0.026$ severe compared with mild sepsis. Thiamin transporter-2: * $p < 0.01$ severe and moderate sepsis compared with control. * $p = 0.005$ mild sepsis compared with control.




Figure 4. Mitochondrial thiamin pyrophosphate transporter (MTPPT) protein (A) and messenger RNA (B). Values are mean \pm SE; $n = 4$ for each group. Representative gel images of MTPPT proteins of different groups were obtained from the same gel. MTPPT protein: ‡ $p = 0.028$ severe sepsis compared with control. ‡ $p = 0.022$ moderate sepsis compared with control. Mitochondrial pyrophosphate transporter mRNA: * $p < 0.001$ severe sepsis compared with control. * $p = 0.004$ moderate sepsis compared with control. ‡ $p = 0.021$ severe compared with mild sepsis.

mortality rate in our study was likely related to the use of systemic antibiotic administered immediately following surgery and maintained for 2 days.

Gut-derived sepsis, the sepsis model employed, has been shown to impair intestinal epithelial barrier function to promote bacterial translocation into the systemic circulation with subsequent bacteremia, and pathologic inflammation of the

intestinal mucosa (11). Whether inflammation of the intestinal mucosa alone with or without adherence of the organisms to the intestinal epithelium is a requirement for thiamin uptake inhibition is unclear. Previous in vitro studies from our laboratory demonstrated that exposure of human-derived intestinal epithelial cells to enteropathogenic *Escherichia coli* (12) or toxin-producing *E. coli* (13) resulted in significant inhibition of

thiamin uptake. Notwithstanding the mechanisms of intestinal thiamin uptake inhibition, the present study demonstrated that sepsis is an important risk factor for the development of thiamin deficiency as previously reported in critically ill patients (2, 4). Sepsis not only inhibited intestinal thiamin uptake and transporters, but the inhibition was a function of sepsis severity. Keeping with previous report in severe septic patients (4), the prevalence of thiamin deficiency was very high at 70%.

Thiamin uptake inhibition in sepsis was also associated with THTR-1 and THTR-2, as well as MTPPT suppression at the protein and mRNA levels (Figs. 2–4). With respect to mRNA suppression, it is unclear at present whether the reduction in mRNA of these transporters is mediated at the transcriptional and/or posttranscriptional levels (e.g., changes in mRNA stability). Further studies are needed to address these issues.

Compared with controls, the significant inhibition of carrier-mediated thiamin uptake in the septic animals was not associated with reduced blood thiamin concentration or thiamin deficiency. This observation was not surprising as our animal model of sepsis was maintained for only 2 days, shorter than thiamin biologic half-life of 9–18 days (14). Additionally, unlike critically ill patients with sepsis whose premorbid conditions may be associated with thiamin deficiency (e.g., chronic alcoholism [15], diabetes [16], celiac disease [17], or long-term use of furosemide diuretic [18]), our animals were healthy prior to the induction of sepsis. Our observation suggests that a normal thiamin blood level in sepsis does not preclude functional thiamin deficiency state, defined as inhibition of carrier-mediated thiamin uptake associated with suppression of THTRs in the cell membrane and mitochondria. Yet, in preexisting thiamin deficiency, the effect of sepsis on carrier-mediated thiamin uptake and its transporters is unknown and requires further study. This is because adaptive regulation to thiamin deficiency occurs. In mice fed with thiamin-deficient diet, carrier-mediated thiamin uptake was upregulated as a result of increased expressions of THTR-2 at both the protein and mRNA levels, without changes in THTR-1 expressions (19). Depending on sepsis severity, it is possible that in preexisting thiamin deficiency, carrier-mediated thiamin uptake and THTR-2 expression may be preserved. However, the septic-induced suppression of THTR-1 as shown in the present study may offset the adaptive regulation of THTR-2.

Sepsis did not only depressed THTR-1 and THTR-2, but also reduced the expression of MTPPT considerably in both the moderate and severe sepsis groups (Fig. 4). In the mitochondria, thiamin pyrophosphate is a cofactor for pyruvate dehydrogenase in the conversion of pyruvate to acetyl-CoA, and α -ketoglutarate dehydrogenase in the conversion of α -ketoglutarate to succinyl-CoA in the tricarboxylic acid cycle (Krebs cycle) for the generation of ATP. Our study showed a significantly reduced ATP generation in animals with moderate and severe sepsis. While sepsis itself, via oxidative stress (11), inhibits electron transport chain complexes activity (20), another pathway for ATP generation independent of thiamin; in the present study, the reduced MTPPT protein or mRNA

expressions contributed to the reduced ATP production. Levels of MTPPT protein and mRNA expressions were responsible for 50% of the variance in ATP production as shown in the significant correlation between ATP production and MTPPT protein and mRNA expression levels (r^2 , 0.50; $p < 0.01$ for MTPPT protein; and r^2 , 0.54; $p < 0.01$ for MTPPT mRNA expression).

ATP is essential to provide energy for numerous cellular processes and reactions (21), particularly in the metabolically active cells such as the intestine for nutrient absorption. As sepsis is a systemic inflammatory syndrome, the reduced ATP generation may not be limited to the intestinal epithelial cells, but may involve other organs via similar mechanism of impaired transport of thiamin pyrophosphate into the mitochondria. Further research is needed to investigate the role of sepsis on functional thiamin deficiency with its consequence of ATP depletion in other organs such as the brain in the manifestation of sepsis-induced delirium (22), or in the peripheral nerves and skeletal muscles in sepsis-induced ICU-acquired muscle weakness.

Limitation of the Study

We defined severity of sepsis according to the site of cecal ligation and not according to the extent of organ failure or tissue hypoperfusion as clinically defined (23). However, animals with severe sepsis had the highest mortality and associated with the highest proinflammatory cytokines levels (Table 1) supporting the severity of sepsis. The duration of sepsis was limited to 2 days. A temporal relationship between duration of sepsis and progression of thiamin deficiency is unknown, and requires further study.

Our measurement of thiamin was obtained from whole blood using microbiological test method rather than using high-performance liquid chromatography as used clinically (24); however, performance of these two methods is comparable (25).

Clinical Implication

Alcoholism has been implicated as the most common predisposing factor for thiamin deficiency with its complications of Wernicke-Korsakoff syndrome and peripheral neuropathy (26). As reported in previous clinical studies (2–5), the results of our study confirmed that sepsis, particularly severe sepsis, is an important risk factor for the development of functional thiamin deficiency via, at least in part, inhibition of carrier-mediated intestinal thiamin uptake. Normal thiamin blood concentration does not preclude functional thiamin deficiency in specific organ. In sepsis, reduced thiamin blood concentration suggests a late manifestation of thiamin deficiency state and may portend a poor prognosis. Further prospective study is needed to confirm this contention.

CONCLUSION

Sepsis induced inhibition of intestinal thiamin uptake; the inhibition was a function of sepsis severity. Sepsis also suppressed THTRs (THTR-1 and THTR-2) and MTPPT, leading to a significant reduction in ATP.

REFERENCES

- Singleton CK, Martin PR: Molecular mechanisms of thiamine utilization. *Curr Mol Med* 2001; 1:197–207
- Donnino MW, Carney E, Cocchi MN, et al: Thiamine deficiency in critically ill patients with sepsis. *J Crit Care* 2010; 25:576–581
- Lima LF, Leite HP, Taddei JA: Low blood thiamine concentrations in children upon admission to the intensive care unit: Risk factors and prognostic significance. *Am J Clin Nutr* 2011; 93:57–61
- Costa NA, Gut AL, de Souza Dorna M, et al: Serum thiamine concentration and oxidative stress as predictors of mortality in patients with septic shock. *J Crit Care* 2014; 29:249–252
- Cruickshank AM, Telfer AB, Shenkin A: Thiamine deficiency in the critically ill. *Intensive Care Med* 1988; 14:384–387
- Said HM: Intestinal absorption of water-soluble vitamins in health and disease. *Biochem J* 2011; 437:357–372
- Rittirsch D, Huber-Lang MS, Flierl MA, et al: Immunodesign of experimental sepsis by cecal ligation and puncture. *Nat Protoc* 2009; 4:31–36
- Chavez-Valdez R, Martin LJ, Flock DL, et al: Necrostatin-1 attenuates mitochondrial dysfunction in neurons and astrocytes following neonatal hypoxia-ischemia. *Neuroscience* 2012; 219:192–203
- Romero-Calvo I, Ocón B, Martínez-Moya P, et al: Reversible Ponceau staining as a loading control alternative to actin in Western blots. *Anal Biochem* 2010; 401:318–320
- Dombrovskiy VY, Martin AA, Sunderram J, et al: Rapid increase in hospitalization and mortality rates for severe sepsis in the United States: A trend analysis from 1993 to 2003. *Crit Care Med* 2007; 35:1244–1250
- Gao M, Jiang Y, Xiao X, et al: Protective effect of pioglitazone on sepsis-induced intestinal injury in a rodent model. *J Surg Res* 2015; 195:550–558
- Ashokkumar B, Kumar JS, Hecht GA, et al: Enteropathogenic *Escherichia coli* inhibits intestinal vitamin B1 (thiamin) uptake: Studies with human-derived intestinal epithelial Caco-2 cells. *Am J Physiol Gastrointest Liver Physiol* 2009; 297:G825–G833
- Ghosal A, Chatterjee NS, Chou T, et al: Enterotoxigenic *Escherichia coli* infection and intestinal thiamin uptake: Studies with intestinal epithelial Caco-2 monolayers. *Am J Physiol Cell Physiol* 2013; 305:C1185–C1191
- Ariaey-Nejad MR, Balaghi M, Baker EM, et al: Thiamin metabolism in man. *Am J Clin Nutr* 1970; 23:764–778
- Tallaksen CM, Bøhmer T, Bell H: Blood and serum thiamin and thiamin phosphate esters concentrations in patients with alcohol dependence syndrome before and after thiamin treatment. *Alcohol Clin Exp Res* 1992; 16:320–325
- Saito N, Kimura M, Kuchiba A, et al: Blood thiamine levels in outpatients with diabetes mellitus. *J Nutr Sci Vitaminol (Tokyo)* 1987; 33:421–430
- Thomson AD: The absorption of radioactive sulphur-labelled thiamine hydrochloride in control subjects and in patients with intestinal malabsorption. *Clin Sci* 1966; 31:167–179
- Seligmann H, Halkin H, Rauchfleisch S, et al: Thiamine deficiency in patients with congestive heart failure receiving long-term furosemide therapy: A pilot study. *Am J Med* 1991; 91:151–155
- Reidling JC, Said HM: Adaptive regulation of intestinal thiamin uptake: Molecular mechanism using wild-type and transgenic mice carrying hTHTR-1 and -2 promoters. *Am J Physiol Gastrointest Liver Physiol* 2005; 288:G1127–G1134
- Peruchi BB, Petronilho F, Rojas HA, et al: Skeletal muscle electron transport chain dysfunction after sepsis in rats. *J Surg Res* 2011; 167:e333–e338
- Stillway LW: The tricarboxylic acid cycle. In: Medical Biochemistry. Fourth Edition. Baynes JW, Dominiczak MH (Eds). Edinburgh, Elsevier, 2014, pp 170–181
- Osiezagha K, Ali S, Freeman C, et al: Thiamine deficiency and delirium. *Innov Clin Neurosci* 2013; 10:26–32
- Dellinger RP, Levy MM, Rhodes A, et al: Surviving Sepsis Campaign Guidelines Committee including the Pediatric Subgroup: Surviving sepsis campaign: International guidelines for management of severe sepsis and septic shock: 2012. *Crit Care Med* 2013; 41:580–637
- Talwar D, Davidson H, Cooney J, et al: Vitamin B(1) status assessed by direct measurement of thiamin pyrophosphate in erythrocytes or whole blood by HPLC: Comparison with erythrocyte transketolase activation assay. *Clin Chem* 2000; 46:704–710
- Sauberlich HE: Newer laboratory methods for assessing nutriture of selected B-complex vitamins. *Annu Rev Nutr* 1984; 4:377–407
- Donnino MW, Vega J, Miller J, et al: Myths and misconceptions of Wernicke's encephalopathy: What every emergency physician should know. *Ann Emerg Med* 2007; 50:715–721